

tients are extremely stoic and fail to react even when the painful area is directly manipulated. Radiographs (X-rays) are usually very helpful in localizing the cause of the lameness.

*Standard radiographic evaluation (x-rays) can provide excellent gross anatomic information in cases of trauma, established arthritis, and advanced bone neoplasia. The changes in bone density necessary to result in identifiable radiographic abnormalities however, preclude early diagnosis of many orthopedic conditions for which scintigraphy is a more sensitive modality.*



Unfortunately many common causes for lameness in dogs and cats do not result in readily demonstrable radiographic lesions until the condition is advanced. This occurs because the structural or anatomic lesion may be extremely small. Fortunately many of these lesions result in a significant change in the metabolism of the bone. This change in the metabolism is what bone scintigraphy is able to detect.

By allowing the detection of the lesion early in the course of the disease, bone scintigraphy may allow for a surgical correction of the problem. Tumors are obviously best treated early in the course of the disease, before they have spread. Conditions causing lameness are usually best treated early in their course, before the development of secondary arthritis which may be incurable.

## Advanced Veterinary Medical Imaging has 2 convenient locations.

### West Los Angeles


Located near the corner of Pico and Sepulveda, conveniently accessible from the 405 and 10 freeways.



2340 Sepulveda Blvd., Los Angeles, CA 90064  
Tel: 310-477-4453

### Orange County

Located in the Jamboree Plaza business park across from the Metrolink Parking lot.



3047 Edinger Ave., Tustin, CA 92780  
Tel: 949-559-7289




## Bone Scintigraphy Update

**Bone Scintigraphy** (or bone scanning) is a valuable tool in the evaluation of the skeletal system. Bone scintigraphy depends on the principal that actively metabolizing bone will incorporate certain bone "tracers." The distribution of these tracers is dependent on the rate of bone turnover and blood flow. The images obtained reveal the relative distribution of these tracers throughout the skeletal system. Since bone scintigraphy reveals changes in bone metabolism more than changes in bone structure, bone scintigraphy complements rather than replaces plain radiography (X-rays). The changes noted on bone scintigraphy usually precede the changes noted on radiographs because the bone metabolism usually changes before the bone structure changes.

The process of bone scintigraphy involves the acquisition of images at various times after the administration of the tracer. The tracer usually requires several hours to be incorporated into the bone and cleared from the remaining tissues of the body. By evaluating the images taken at different times after the administration of the tracer, information about blood flow versus soft tissue (muscle, ligament, tendon) and bone metabolism can be obtained.

Bone scintigraphy is useful in the evaluation of the cancer patient with a primary bone tumor or a tumor of other than bone origin suspected of possible metastasis, or spread to the skeletal system. Bone scintigraphy is also useful in the evaluation of osteomyelitis or infected bone. This condition can occur independent of other disease or as a complication to previous injuries or fracture repair.

One of the most useful applications for bone scintigraphy is in the evaluation of the patient with a poorly localized lameness. Localization of the source of a lameness in a veterinary patient is frequently difficult. This difficulty is partially due to the inability of our patients to communicate the source of their pain and is complicated by the fact that many of our pa-




*Figure 1. Multiple delayed images of a dog with a primary osteosarcoma (bone tumor) in the left skull with diffuse metastatic disease.*



*Figure 2. Soft tissue and delayed bone images of a 5 year old German Shepherd dog with a right rear leg lameness due to an avulsion of the cranial cruciate ligament. This image shows a very focal lesion in the bone at the origin of the cranial cruciate ligament. Radiographs were normal. The dog responded well to surgical repair of the cruciate ligament.*

Any condition which results in a change in the metabolism of the bone will result in a change in the appearance of the bone scan. As a result lesions like fractures, infections, tumors, and arthritis can be recognized with a bone scan long before they can be seen with plain radiographs.



*Figure 3. Three phase bone scintigraphy demonstrating a typical pattern for fragmented medial coronoid disease (FCP). Notice the subtle increase in blood flow to the left elbow, mildly increased radionuclide distribution to the left elbow on the soft tissue phase and focal area of increased radionuclide uptake in the left elbow on the delayed bone images.*